



## Use of GLP-1 mimetic in type 2 diabetes mellitus: is it the end of fragility fractures?

Submitted by Guillaume Mabillean on Wed, 11/26/2014 - 17:24

Titre	Use of GLP-1 mimetic in type 2 diabetes mellitus: is it the end of fragility fractures?
Type de publication	Article de revue
Auteur	Mabillean, Guillaume [1]
Editeur	Springer-Verlag
Type	Article scientifique dans une revue à comité de lecture
Année	2015
Langue	Anglais
Date	2014 Oct 28
Pagination	1-2
Volume	48
Section	1
Titre de la revue	Endocrine
ISSN	1559-0100
Résumé en anglais	Type 2 diabetes mellitus (T2DM) is an insidious disease characterized by hyperglycemia and hyperinsulinemia. Among all complications of T2DM, bone fractures represent a severe alteration of the quality of life of affected individuals and are cost-effective. One explanation for bone fracture could be linked to other diabetes complications such as retinopathy and neuropathy, which increase the incidence of falls. Another explanation could reside in a direct effect of T2DM on bone resulting in poor "quality" of the bone matrix.[...]
URL de la notice	<a href="http://okina.univ-angers.fr/publications/ua5554">http://okina.univ-angers.fr/publications/ua5554</a> [2]
DOI	<a href="https://doi.org/10.1007/s12020-014-0461-1">10.1007/s12020-014-0461-1</a> [3]
Autre titre	Endocrine
Identifiant (ID) PubMed	25348148 [4]

---

### Liens

[1] <http://okina.univ-angers.fr/guillaume.mabillean/publications>

[2] <http://okina.univ-angers.fr/publications/ua5554>

[3] <http://dx.doi.org/10.1007/s12020-014-0461-1>

[4] <http://www.ncbi.nlm.nih.gov/pubmed/25348148?dopt=Abstract>

Publié sur *Okina* (<http://okina.univ-angers.fr>)